



Data  
Models  
Inventories

# PARIS

Process Attribution of Regional Emissions

GA 101081430, RIA

Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

---

## Deliverable 4.3

---

Delivery due date Annex I	Month 28 - 01/06/2025
Actual date of submission	01/02/2026
Lead beneficiary: UEDIN	Work package: 4    Nature:    Dissemination level: Public Report/data
Responsible scientist	Tim Arnold
Contributors	Stephan Henne (EMPA), Anita Ganesan UNIVBRIS, Alistair Manning MO
Internal reviewers	
Version: 0.1	

---



Horizon Europe Cluster 5: Climate, energy and mobility

*"This project has received funding from the European Union's Horizon Europe Research and Innovation programme under HORIZON-CL5-2022-D1-02 Grant Agreement No 101081430 - PARIS".*

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

## Table of content

<b>1. CHANGES WITH RESPECT TO THE DOA (DESCRIPTION OF THE ACTION)</b>	<b>3</b>
<b>2. DISSEMINATION AND UPTAKE</b>	<b>3</b>
<b>3. SHORT SUMMARY OF RESULTS</b>	<b>3</b>
<b>4. EVIDENCE OF ACCOMPLISHMENT</b>	<b>3</b>
4.1 INTRODUCTION   BACKGROUND OF THE DELIVERABLE	3
4.2 SCOPE OF THE DELIVERABLE   MILESTONE	4
4.3 CONTENT OF THE DELIVERABLE   MILESTONE	4
4.4 CONCLUSION AND POSSIBLE IMPACT	10
4.5 REFERENCES	10
<b>5. HISTORY OF THE DOCUMENT</b>	<b>10</b>

## 1. Changes with respect to the DoA (Description of the Action)

Although the report and associated data upload were delayed beyond the original timeline, the contents of the deliverables and milestone outcomes are unchanged from the DoA.

## 2. Dissemination and uptake

The data uploaded to the ICOS Carbon Portal form the basis for Annex reports in WP2. As such they have been shared with the relevant project partners prior to the completion of D2.3. Final data products are all available publicly on the ICOS Carbon Portal for use by other scientists and practitioners and have been shared with national inventory compilers (e.g., the Dutch inventory team has based the figures produced by themselves for their annex to the 2026 NID report on these data).

## 3. Short Summary of results

The inversion modelling systems employed in PARIS, represented by a total of six combinations of inversion systems (RHIME, InTEM, ELRIS) and transport models (NAME, FLEXPART) were used to estimate CH<sub>4</sub> fluxes across Europe for the common period 2016 to 2024, with NAME-based inversions starting earlier. Inverse modelling results include monthly CH<sub>4</sub> fluxes at a spatial resolution of 25 km x 25 km and country aggregates for the European domain. These were collected in commonly defined data formats and made publicly available (<https://doi.org/10.18160/GR1Q-6SK4>). Results were visualised by the inversion-intercomparison tool fluxy and figures of the included data were semi-automatically produced, forming the basis of the PARIS annex drafts submitted to the national inventory teams of the PARIS focus countries.

## 4. Evidence of accomplishment

Inverse modelling results (including flux estimates as well as observed and simulated atmospheric mole fractions) from the individual inversions are available on the ICOS Carbon Portal:

ELRIS-FLEXPART, <https://meta.icos-cp.eu/objects/od9o-Uka9A0b1IV-Z0lk-WeF>

ELRIS-NAME, <https://meta.icos-cp.eu/objects/gp1oRDdSjjeWRC805PY6-zFH>

InTEM-FLEXPART, [https://meta.icos-cp.eu/objects/5W5IE8X5xTyslvq\\_FjrRv-kV](https://meta.icos-cp.eu/objects/5W5IE8X5xTyslvq_FjrRv-kV)

InTEM-NAME, [https://meta.icos-cp.eu/objects/TFCu1I52WEBjnW\\_2jwaDhb0o](https://meta.icos-cp.eu/objects/TFCu1I52WEBjnW_2jwaDhb0o)

RHIME-FLEXPART, [https://meta.icos-cp.eu/objects/b4lh\\_958ogPzX70cv-YrvZ4Y](https://meta.icos-cp.eu/objects/b4lh_958ogPzX70cv-YrvZ4Y)

RHIME-NAME, [https://meta.icos-cp.eu/objects/XIH2WO6haVE3HdoRvApr\\_5yy](https://meta.icos-cp.eu/objects/XIH2WO6haVE3HdoRvApr_5yy)

### 4.1 Introduction | Background of the deliverable

Inverse model estimates based on atmospheric observations are one of the central results of the PARIS projects as these are shared with national inventory teams to validate their bottom-up reporting. The observation-based inverse estimate is considered to be largely independent of the bottom-up reporting and allows the identifications of shortcomings in the bottom-up reporting in areas well covered by the observing network. In the case of

#### D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

PARIS eight focus countries were chosen for inter-comparison, most of which exhibit good long-term observational coverage, with few targeted because of emerging new observations. The primary result of the inversion systems applied in PARIS are spatially resolved fluxes with, in the case of CH<sub>4</sub>, monthly resolution. These data are aggregated to national totals on timescales (annual) relevant for GHG reporting to UNFCCC. Hence, an actionable dataset is created for direct comparison for the inventory teams. Inversion results are delivered in a timely manner, meaning that the results presented here and communicated to inventory teams in October 2025, contain emission estimates up to the end of 2024. In April 2026 the national inventories need to include emission estimates for 2024 as well. Therefore, the PARIS dataset allows direct comparison up to the latest possible reporting year. Details of the comparison of inversion results and UNFCCC reporting are contained in PARIS deliverable D2.3. Here, the delivered dataset and its use is outlined.

#### 4.2 Scope of the deliverable | milestone

The delivered dataset comprises inverse modelling results of European CH<sub>4</sub> surface fluxes from six different combinations of inverse modelling systems and transport models. The common period 2016-2024 is covered for which the observational network is most dense, and uncertainties of flux estimates smallest.

#### 4.3 Content of the deliverable | milestone

The inverse methodology was described in detail elsewhere (e.g., model documentation supplement to D2.3). Performance of the chosen inverse setup for CH<sub>4</sub> was analysed in M39. Here, the data product itself is described, whereas interpretation of the inverse modelling results is provided through D2.3.

The uploaded datasets comprise three types of data:

1. Monthly CH<sub>4</sub> flux estimates (including prior and posterior estimates and their uncertainties) at a spatial resolution of 25 km x 25 km  
 Relevant variables are described as part of the standardised netcdf format for inversion results (flux file) in **Error! Reference source not found.** and [https://github.com/openghg/fluxy/blob/devel/data/templates/README\\_templates.md](https://github.com/openghg/fluxy/blob/devel/data/templates/README_templates.md)
2. Country aggregates of monthly CH<sub>4</sub> fluxes for western European countries (areas well-covered by observations network). Prior and posterior values and their uncertainties and covariances between individual regions are reported. These data are contained in the same files the spatially resolved fluxes.  
 Relevant variables are described as part of the standardised netcdf format for inversion results in **Error! Reference source not found.** and (flux file): [https://github.com/openghg/fluxy/blob/devel/data/templates/README\\_templates.md](https://github.com/openghg/fluxy/blob/devel/data/templates/README_templates.md)
3. Time series of observed and simulated CH<sub>4</sub> mole fractions

#### D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

Relevant variables are described as part of the standardised netcdf format for inversion results in *Table* and (concentration file):

[https://github.com/openghg/fluxy/blob/devel/data/templates/README\\_templates.md](https://github.com/openghg/fluxy/blob/devel/data/templates/README_templates.md)

Each of the six sensitivity inversions that form the PARIS ensemble is available as a pair of flux and concentration file. In the following a few example plots of the available data are presented as created with the comparison tool fluxy.

Spatially resolved flux information is shown in Fig. 1 for all six PARIS inversions. The displayed fluxes are the a priori (netcdf variable `flux_total_prior`) and a posteriori fluxes (`flux_total_posterior`) averaged from the monthly data contained in the dataset to a mean for the period January 2016 to December 2024. Since the employed inversion systems all work with a reduced inversion grid (regions in which individual factors are optimised to arrive at the a posteriori results), two additional variables are provided in the netcdf files which provide results at the resolution of these reduced inversion grids (`flux_total_posterior_inversion_grid`, `flux_total_prior_inversion_grid`). Furthermore, uncertainties of the spatially resolved fluxes are provided in variables `stdev_flux_total_prior` and `stdev_flux_total_posterior` or, if non-Gaussian uncertainties are involved, as a percentile range in variables `percentile_flux_total_prior` and `percentile_flux_total_posterior`.

An example of country total CH<sub>4</sub> flux data is given in Fig. 2. The displayed fluxes are the a priori (`flux_total_prior_country`) and a posteriori estimates (`flux_total_posterior_country`), available at monthly resolution and collected across all six inversion datasets. Uncertainty bands for the a posteriori either are given either by the standard deviation of a Gaussian distribution (`stdev_total_posterior_country`) or by a percentile range for non-Gaussian distributions (`percentile_total_posterior_country`). The uncertainty of the a priori fluxes is contained dataset (`stdev_total_prior_country`, `percentile_total_prior_country`) but omitted from the plot. Not contained in the dataset are the countries' reported emissions (black bars). These need to be obtained from the common reporting tables (CRT) available at UNFCCC.

Time series of observed and simulated CH<sub>4</sub> mole fractions are discussed as part of M41 and various visualisations can be found therein. Variables are listed in Table 3. Several of these variables are optional because different inversion systems make different assumptions about how to split the total mole fraction (at least a split into boundary conditions (`mf_bc_prior` and `mf_bc_posterior`) and total mole fractions (`mf_prior` and `mf_posterior`) is available).

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

**Table 1:** Variables for spatially resolved fluxes defined in the standardised netcdf format for inversion results (flux file).

Grid variables	Units	Dimensions	Description
flux_total_prior	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Prior total <species> fluxes
flux_total_posterior	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Posterior total <species> fluxes
stdev_flux_total_prior	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Standard deviation of prior total <species> fluxes
stdev_flux_total_posterior	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Standard deviation of posterior total <species> fluxes
<b>Optional variables</b>			
flux_total_prior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Prior total <species> fluxes on the reduced inversion grid
flux_total_posterior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Posterior total <species> fluxes on the reduced inversion grid
stdev_flux_total_prior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Standard deviation of prior total <species> fluxes on the reduced inversion grid
stdev_flux_total_posterior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, latitude, longitude	Standard deviation of posterior total <species> fluxes on the reduced inversion grid
<b>Alternative to stdev for non-Gaussian PDFs</b>			
percentile_flux_total_prior	mol m <sup>-2</sup> s <sup>-1</sup>	time, percentile, latitude, longitude	Percentile of prior total <species> fluxes
percentile_flux_total_posterior	mol m <sup>-2</sup> s <sup>-1</sup>	time, percentile, latitude, longitude	Percentile of posterior total <species> fluxes
percentile_flux_total_prior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, percentile, latitude, longitude	Percentile of prior total <species> fluxes on the reduced inversion grid
percentile_flux_total_posterior_inversion_grid	mol m <sup>-2</sup> s <sup>-1</sup>	time, percentile, latitude, longitude	Percentile of posterior total <species> fluxes on the reduced inversion grid

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

**Table 2:** Variables for country-aggregated fluxes defined in the standardised netcdf format for inversion results (flux file).


By-country variables	Units	Dimensions	Description
flux_total_prior_country	kg yr <sup>-1</sup>	time, country	Country-total prior <species> fluxes
flux_total_posterior_country	kg yr <sup>-1</sup>	time, country	Country-total posterior <species> fluxes
stdev_flux_total_prior_country	kg yr <sup>-1</sup>	time, country	Standard deviation of country-total prior <species> fluxes
stdev_flux_total_posterior_country	kg yr <sup>-1</sup>	time, country	Standard deviation of country-total posterior <species> fluxes
<b>Alternative to stdev for non-Gaussian PDFs</b>			
percentile_flux_total_prior_country	kg yr <sup>-1</sup>	time, percentile, country	Percentiles of country-total prior <species> fluxes
percentile_flux_total_posterior_country	kg yr <sup>-1</sup>	time, percentile, country	Percentiles of country-total posterior <species> fluxes
<b>Optional</b>			
covariance_flux_total_posterior_country	kg <sup>2</sup> yr <sup>-2</sup>	time, country, country	Covariance of country-total posterior <species> fluxes

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round


**Table 3:** Variables defined for observed and simulated concentrations at observing sites included in the standardised netcdf format for inversion results (concentration file).

Observation variables	Units	Dimensions	Description
platform	-	platform	Identifier of observing platform
mf_observed	mol mol <sup>-1</sup>	index	Observed mole fraction of <species> in dry air
<b>Simulated variables</b>			
mf_prior	mol mol <sup>-1</sup>	index	Prior simulated mole fraction of <species> in dry air
mf_posterior	mol mol <sup>-1</sup>	index	Posterior simulated mole fraction of <species> in dry air
mf_bc_prior	mol mol <sup>-1</sup>	index	Prior simulated boundary condition mole fraction including site bias
mf_bc_posterior	mol mol <sup>-1</sup>	index	Posterior simulated boundary condition mole fraction including site bias
<b>Optional</b>			
stdev_mf_prior	mol mol <sup>-1</sup>	index	Standard deviation of prior simulated mole fractions due to state vector uncertainty
stdev_mf_posterior	mol mol <sup>-1</sup>	index	Standard deviation of posterior simulated mole fractions due to state vector uncertainty
mf_bias_prior	mol mol <sup>-1</sup>	index	Prior simulated mole fraction site bias
mf_bias_posterior	mol mol <sup>-1</sup>	index	Posterior simulated mole fraction site bias
mf_outer_prior	mol mol <sup>-1</sup>	index	Prior simulated mole fraction contribution from distant regions
mf_outer_posterior	mol mol <sup>-1</sup>	index	Posterior simulated mole fraction contribution from distant regions
<b>Alternative to stdev for non-Gaussian PDFs</b>			
percentile_mf_prior	mol mol <sup>-1</sup>	index, percentile	Percentile of prior simulated mole fraction due to state vector uncertainty
percentile_mf_posterior	mol mol <sup>-1</sup>	index, percentile	Percentile of posterior simulated mole fraction due to state vector uncertainty
<b>Uncertainty variables</b>			
stdev_mf_total	mol mol <sup>-1</sup>	index	Total model-data-mismatch uncertainty applied in inversion
<b>Optional</b>			
stdev_mf_observed_repeatability	mol mol <sup>-1</sup>	index	Repeatability uncertainty of observed mole fraction
stdev_mf_observed_variability	mol mol <sup>-1</sup>	index	Variability of observed mole fraction within aggregation interval
stdev_mf_model	mol mol <sup>-1</sup>	index	Model uncertainty of simulated mole fraction

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round



**Fig. 1:** A priori (top), a posteriori (middle) fluxes and their differences (bottom) for the six PARIS CH<sub>4</sub> inversions (columns). For each panel the monthly flux data was aggregated for the period January 2016 to December 2024.



**Fig. 2:** Time series of country total, monthly a priori (dashed lines), a posteriori (solid lines) CH<sub>4</sub> emissions and their uncertainties (ribbons). For comparison, bottom-up (inventory data not part of this deliverable) flux estimates are shown with annual bars.

D4.3 - Full year of CH<sub>4</sub> data, uploaded to the ICOS portal - 1<sup>st</sup> round

#### 4.4 Conclusion and possible impact

The uploaded inversion results form the basis for the production of PARIS draft annexes (D2.3) and are shared with the national inventory teams and wider scientific community. Results will be directly used by the Dutch and Irish inventory compilers in their upcoming (2026) inventory document for the UNFCCC. An update of the current dataset is expected by October 2026 and will then include flux estimates until the end of 2025.

The standardised data format and the open-source fluxy visualisation and inter-comparison tool (<https://github.com/openghg/fluxy>) ensure easy access to the inversion results and inter-operability with other inverse modelling groups)

#### 4.5 References

Ganesan, A., Manning, A., Henne, S., De Longueville, H., Ramsden, A., Brito Melo, D., Danjou, A., Andrews, P., Murphy, B., Redington, A., and Pitt, J.: Inverse modelling results for European non-CO<sub>2</sub> greenhouse gas emissions, ICOS ERIC - Carbon Portal, 2026.

### 5. History of the document

Version	Author(s)	Date	Changes
0.1	Tim Arnold	2026-02-19	First draft